The Guide to Community Preventive Services - **TOBACCO**

Patricia A. Nolan, MD, MPH Congressional Prevention Caucus Briefing – June 8, 2007

The Community Guide

- > A Task Force
- > A Process
 - Systematic reviews of the literature
 - Assessments of study quality
 - Summary of outcomes
- > A Product
 - Evidence-based recommendations
 - Book and the web site

Methods

- > Develop a conceptual framework
- > Search for evidence
- > Rate the quality of the evidence
- > Summarize the information
- Translate the strength of evidence into recommendations for use

Barriers Intervention Intended Outcomes Additional Outcomes Benefits Reduced Morbidity and/or Mortality

Let's Talk About Evidence

- > Sufficient and Insufficient Evidence
 - The number and quality of studies
 - The clarity of the causal relationships
- > Strong and Weak Evidence
 - The size of the effects
 - The consistency of the effects
- > Program Examples
 - · Real world undertakings based on evidence

Using the Evidence for Community Interventions -- TOBACCO

- Health care interventions are directed at helping individuals
 - Those with tobacco-related disease
 - Those who want to quit
- > Community interventions are directed toward -
 - Decreasing tobacco use
 - Decreasing the number of tobacco users
 - Decreasing exposure to environmental tobacco smoke

At the community level ...

Decrease smoking initiation by children, youth and young adults by

- Increasing the unit price of tobacco products
 Recommended on the basis of strong evidence
- Using mass media campaigns in combination with other interventions
 - Recommended on the basis of strong evidence

At the community level ...

Increase smoking cessation by

- Increasing the unit price of cigarettes Recommended on the basis of strong evidence
- Using mass media campaigns to stimulate and support smoking cessation
 - Recommended on the basis of strong evidence

In the Health Care System...

Increase smoking cessation by

 Using provider reminder systems and provider education to encourage smoking cessation interventions

Recommended on the basis of strong evidence

> Providing telephone support (quit lines) in combination with other interventions

Recommended on the basis of strong evidence

In the Health Care System...

Increase smoking cessation by

 Reducing out-of-pocket costs of effective clinical cessation interventions

Recommended on the basis of sufficient evidence

Using provider prompts to encourage smoking cessation intervention

Recommended on the basis of sufficient evidence

At the community level ...

Decrease exposure to tobacco smoke in the environment by

- Smoking bans and policies
 Recommended on the basis of strong evidence
- > No harms to health were identified

At the Community Level...

Choosing community interventions

- Shared experiences from state and local health departments and from community agencies
- Research on effective interventions in special populations and sites
- Interaction among community and medical interventions.

The Guide to Community Preventive Services - TOBACCO

A great source for evidence :

www.thecommunityguide.org

Thanks to

- The Community Guide Task Force members
- > The Community Guide staff who sustain this effort
- > Dr. David Hopkins for several of these slides
- > To you for your interest in eliminating the scourge of tobacco